

# Techniques frequently used during London Olympic judo tournaments: A biomechanical approach

## Authors' Contribution:

- A** Study Design
- B** Data Collection
- C** Statistical Analysis
- D** Manuscript Preparation
- E** Funds Collection

**Stanislaw Sterkowicz<sup>1ABCDE</sup>, Attilio Sacripanti<sup>2BCDE</sup>, Katarzyna Sterkowicz-Przybycień<sup>3ABCDE</sup>**

<sup>1</sup> Department of Theory of Sport and Kinesiology, Institute of Sport, University School of Physical Education, Kraków, Poland

<sup>2</sup> Chair of Biomechanics of Sports, FIJKAM, ENEA, University of Rome "Tor Vergata", Italy

<sup>3</sup> Department of Gymnastics, Institute of Sport, University School of Physical Education, Kraków, Poland

**Source of support:** Departmental sources

**Received:** 17 December 2012; **Accepted:** 18 March 2013; **Published online:** 25 March 2013

**ICID:** 883848

## Abstract

### Background & Study Aim:

Feedback between training and competition should be considered in athletic training. The aim of the study was contemporary coaching tendencies in women's and men's judo with particular focus on a biomechanical classification of throws and grappling actions.

### Material & Methods:

359 throws and 77 grappling techniques scored by male and female athletes in Olympic Judo Tournaments (London 2012) have been analysed. Independence of traits (gender and weight category by technique classes) was verified via  $\chi^2$  test. Comparison between frequency of each subsequent technique class and rest/inconclusive counts was made in 2x2 contingency tables. The significance level was set at  $p \leq 0.05$ .

### Results:

Throwing technique frequencies grouped in the seven biomechanical classes were dependent on gender. A significant difference was found between frequencies of variable arm of physical lever technique scored by males (27.09%) and females (16.67%) as compared to the rest/inconclusively techniques counts. Significant differences between men who competed in extra lightweight and heavy weight concerned the frequency of the techniques used with maximum arm or variable arm of physical lever and a couple of forces applied by trunk and legs. In females, a tendency to higher frequency of techniques that used couple of forces applied by arm or arms and leg was observed in extra lightweight compared to the heavy weight.

### Conclusions:

Because the technique preferred in the fight depends on a gender and weight category of a judoka, the relationships found in this study, which can be justified by the biomechanics of throws, should be taken into consideration in technical and tactical coaching of the contestants. A method used in this study can be recommended for future research concerning coaching tendencies.

### Key words:

judo • biomechanical classification • technical analysis

### Author's address:

Attilio Sacripanti, ENEA – Casaccia Research Centre, Via Anguillarese 301, 00060 Rome, Italy;  
e-mail: attilio.sacripanti@uniroma2.it

## BACKGROUND

Observation of the fight in judo competitions is necessary as it is the only opportunity for verification of the process of the contestant's coaching. This feedback is particularly important if new rivals emerge in a weight category or if fighting regulations force athletes and coaches to face new, more demanding challenges [1].

World championships and the Olympic Games are the moments that best summarize many years of training. Male [2,3] and, more recently, female [4] competitions have been analysed in detail during these events, both in groups and individually [4]. These analyses used a traditional classification of techniques developed in the Kodokan Judo Institute [5], with throws including hand techniques, loin techniques, foot and leg techniques,

**GAI** (General Action Invariants) – all the trajectories applied to shorten the distance between athletes [10].

**(SAI)** – “Specific Action Invariants”, which can be split into Superior Specific Action Invariants (SSAI) and Inferior Specific Action Invariants (ISAI) all the movements performed by Athlete’s kinetic chains [10].

**“Innovative Throws”** – are all throwing techniques that keep alive the formal aspect of classic judo throws, and differ in terms of grips and final direction of applied forces only [12].

**“New or Chaotic Throws”** – principally belong to the lever-type Group, and are characterized by the application of different GAI trajectory, grips positions (SSAI) which apply force in different (non-traditional) directions while simultaneously applying (ISAI) stopping point in non-classical positions [10].

and the art of throwing in a horizontal posture, sacrificing his own body’s balance. In addition, the art of grappling encompasses holds, strangling, and locking the joints (elbows). The need for improvement of technical and tactical preparation of contestants has caused that some additional criteria of joint classification have also been used in practice: based on the direction of *kuzushi* (the action taken by one contestant in order to throw the other contestant out of balance), presence of body rotation performed by a thrower and tactical situation when performing a throw i.e. a single attack, combination or counterattack [6]. The observations and analysis of the course of the fight during competitions at the elite level have been the focus of studies that yielded results that are useful for both theory and practice of judo and judo coaching. At the same time, a consistent throw classification was developed based on the biomechanical criteria [7,8], which has not been used for the analysis of the frequency of actions in a judo fight so far.

### Concept of the work

All judo athletes used the same fighting rules regardless of whether they were males or females. It was assumed that the frequency found for a particular technique class might be related to gender or a weight category. Therefore, it seemed justified to formulate the following research questions: (1) Which throw techniques and grappling actions (Sacripanti’s biomechanical criterion) are frequently used by contemporary elite judo athletes?; (2) Were there any differences between the frequency observed in male and female competitors?; (3) Were there any differences depending on weight categories?; (4) How often did the penalties for non-combativity in judo fights occur? The aim of the present study was contemporary coaching tendencies in women’s and men’s judo with particular focus on biomechanical classification of throws.

### MATERIAL AND METHODS

There are no ethical issues involved in the analysis and interpretation of the data used as these were obtained from other sources and were not generated by experimentation. The athletes’ personal identification was replaced by a code, which ensured anonymity and confidentiality. All actions of male and female athletes were recorded using IJF coding system [9]. There were 359 judo-throw-techniques and 77 grappling actions performed and scored during Olympic tournaments in London. Penalties caused by the breach of judo fight regulations (n=591) were also analysed. Furthermore, each technique was rearranged by us into the biomechanical classification system [7,8,10]. A data analysis was conducted for identification of each technique

within nine classification groups (Tables 4A and 4B in Annex [8,11,12]). Frequency of technique count distribution was compared using Statgraphics Centurion XVI.I software. Independent variables were gender (males; females) and weight categories: (1) extra lightweight; (2) half lightweight, lightweight, half middleweight; (3) middleweight; half heavyweight; (4) heavy weight. The rationale behind this division was separation of semi-open categories i.e. 1<sup>st</sup> group (upper limits, only) and 4<sup>th</sup> group (lower limits, only). The ranges between the limits in the 2<sup>nd</sup> group were 21.0% and 19.0% of the lower limit of heavy weight of men and female. In the 3<sup>rd</sup> group, these ranges were similar (19.0% and 19.2%, respectively). In the multi-way tables, due to the expectedly small numbers, independence of traits was verified with  $\chi^2$  test in the logarithmic form (G-test) [13]. Comparisons between the frequency of each subsequent technique classes and a rest counts were presented in 2×2 contingency tables. The Yates correction verified by Fisher exact test (FET) was used for small data. In the case of the significant dependency, the contingency coefficient C was calculated. The significance level was set at  $p \leq 0.05$ .

### RESULTS

Table 1 presents the frequency of techniques used by males and females from different weight categories. In general, techniques based on a couple of forces were used less frequently (39.6%) than the techniques used with physical lever (60.5%). There were no significant differences ( $\chi^2=0.875$ ,  $df=1$ ,  $p=0.350$ ) between the frequencies of these techniques performed by males (37.4% vs. 62.6%) and females (42.3% vs. 57.7%). Technique frequencies grouped in the seven biomechanical classes (Table 1) were dependent on gender ( $\chi^2=16.00$ ,  $df=6$ ,  $p<0.05$ ,  $C=0.207$ ).

The techniques used with maximum physical lever were scored the most often (25.1%), independently of the competitor’s gender ( $p>0.05$ ), i.e. 24.1% in male and 26.3% in female judokas. In those techniques, the group of *tai-otoshi* was a typical throw scored (7.0%) with similar frequencies in males and females (6.4% and 7.7%, respectively). Using the biomechanical criterion, the next frequently scored throws were those using a variable arm of physical lever (22.6%). Significant differences ( $\chi^2=5.24$ ,  $df=1$ ,  $p=0.022$ ,  $C=0.120$ ) were found between the frequencies of this class of techniques scored by males (27.1%) and females (16.7%) as compared to the rest/inconclusively performed techniques count. *Seoi-nage* is an example of throw (14.8%) which was less frequently used by females (8.3%) compared to males (19.7%). The frequencies of other techniques classified within next five classification groups

**Table 1.** Frequency of throw techniques used during Judo Olympic Tournaments (2012) by males and females from different weight categories.

Technique of throws codes CODE	Total	Males	Females	Male groups/weight categories				Female groups/weight categories			
				Group 1	Group 2	Group 3	Group 4	Group 1	Group 2	Group 3	Group 4
PLmaxA	90	49	41	9	24	14	2 <sup>#</sup>	8	14	12	7
PLvA	81	55	26*	17	23**	14**	1**	2	13	8	3
CAL	77	40	37	9	17	11	3	6	15	15 <sup>#</sup>	1 <sup>#</sup>
CTL	55	27	28	2	12	8	5*	1	12	12	3
PLminA	35	20	15	2	7	8	3	1	7	3	4
PLmidA	11	3	8	0	1	1	1	0	4	2	2
CA	10	9	1	0	6	2	1	0	0	0	1
Total	359	203	156	39	90	58	16	18	65	52	21

PLmaxA – Physical lever applied with max arm; PLvA – Physical lever applied with variable arm; CAL – Couple of forces applied by arm or arms and leg; CTL – Couple of forces applied by trunk and legs; PLminA – Physical lever applied with min arm; PLmidA – Physical lever applied with middle arm; CA – Couple of forces applied by arms. \* Significant difference between males and females; # significant difference between group 1 and group 4; \*\* significant difference between group 1 and 2, 3, 4 groups, ## significant difference between groups 2<sup>nd</sup> and 3<sup>rd</sup>.

(see Table 1) did not differ significantly between females and males ( $p>0.05$ ). In the hierarchy of frequently used techniques, the subsequent places were ranked: 3. techniques of couple forces applied by arm(s) and leg (21.45%) within representative *o-uchi-gari* and *ko-uchi-gari* (6.1% each). 4. Techniques of couple forces applied by trunk and legs, with typical technique being *uchi-mata* (7.0%; but 4.9% in males and 9.6% in females) 5. Techniques applied by minimum physical lever (9.75%), were typically *soto-makikomi* (3.0% in males and 5.8% in females) 6. Techniques applied by medium physical lever (3.06%) and 7. Techniques of couple forces applied by arms (2.79%). Techniques of couple forces applied by trunk and arms only (like *morote-gari*) and legs only (like *kani-basami*) were not observed because they were not allowed by the regulations for contemporary official competitions.

In males, the frequency distribution of the PLmaxA techniques and rest/inconclusive techniques performed by competitors from half-open weight categories (1<sup>st</sup> vs. 4<sup>th</sup>) differed significantly ( $\chi^2=5.24$ ,  $df=1$ ,  $p<0.05$ ,  $C=0.294$ ). A comparison of the frequency of using PLvA techniques to all other techniques shows that the athletes from the group 1<sup>st</sup> used them significantly more often than the 2<sup>nd</sup> group ( $\chi^2=4.137$ ,  $p=0.042$ ,  $C=0.176$ ), the 3<sup>rd</sup> group ( $\chi^2=4.058$ ,  $p=0.044$ ,  $C=0.200$ ), and the 4<sup>th</sup> group ( $\chi^2=8.642$ ,  $df=1$ ,  $p<0.004$ ,  $C=0.368$ ). Weight categories and CAL technique frequencies were independent at 95% confidence level ( $p>0.05$ ). The competitors from the group 4<sup>th</sup> performed CTL techniques ( $\chi^2=4.816$ ,  $df=1$ ,  $p=0.028$ ,  $C=0.284$ ) more efficiently

than group 1<sup>st</sup>. Other results for male-specific techniques compared to the inconclusive group were insignificant.

A comparison between a particular technique and the rest/inconclusive techniques in females shows a tendency to higher frequency of CAL technique used in 1<sup>st</sup> compared to the 4<sup>th</sup> group ( $\chi^2=3.608$ ,  $df=1$ ,  $p=0.057$ ,  $C=0.291$ ;  $FET=0.035$ ). The CAL technique was also relatively often used in 3<sup>rd</sup> compared to the 4<sup>th</sup> weight category ( $\chi^2=3.760$ ,  $df=1$ ,  $p=0.053$ ,  $C=0.176$ ;  $FET=0.029$ ). In gripping actions, women lost because of the vascular chokes more often than men (Table 2). Men from the 1<sup>st</sup> weight category performed pinning techniques of the four corner type ( $\chi^2=4.024$ ,  $df=1$ ,  $p=0.045$ ,  $C=0.191$ ) much less frequently than those from the 3<sup>rd</sup> category. Among women, the number of particular grappling techniques did not depend on weight categories.

The frequency of penalties for non-combativity was significantly higher among men (65.4%) than in women (55.5%, cell's percentage of the column) ( $\chi^2=9.783$ ,  $df=1$ ,  $p=0.002$ ,  $C=0.128$ ). The men from extra light-weight category were imposed penalties for non-combativity less frequently (56.8%) than those from the heavy weight (83.9%) ( $\chi^2=8.797$ ,  $df=1$ ,  $p=0.003$ ,  $C=0.286$ ). Similar results were observed for comparison of the 2<sup>nd</sup> (58.3%) and 4<sup>th</sup> groups (83.9%) ( $\chi^2=12.985$ ,  $df=1$ ,  $p<0.001$ ,  $C=0.236$ ) as well as the 3<sup>rd</sup> (68.6%) and 4<sup>th</sup> groups (83.9%) ( $\chi^2=4.703$ ,  $df=1$ ,  $p<0.030$ ,  $C=0.167$ ). No relationships were found between the frequency of penalties and weight category in women (Table 3).

**Table 2.** Frequency of grappling techniques used during Judo Olympic Tournaments (2012) by males and females from different weight categories.

Grappling actions codes	Total	Males	Females	Male groups/weight categories				Female groups/weight categories			
				Group 1	Group 2	Group 3	Group 4	Group 1	Group 2	Group 3	Group 4
KGARAMI	1	0	1	0	0	0	0	0	0	0	1
KHISHIGI	19	9	10	5	3	1	0	0	6	3	1
OKESA	11	7	4	2	2	1	2	0	3	1	0
OSHIHO	30	19	11	2	9	6 <sup>#</sup>	2	1	2	5	3
SRESP	12	8	4	2	4	0	2	1	3	0	0
SVASC	4	0	4 <sup>*</sup>	0	0	0	0	2	2	0	0
Total	77	43	34	11	18	8	6	4	16	9	5

KGARAMI – joint techniques of the entangled joint lock type; KHISHIGI – joint techniques of the bending and pressing against elbow joint type; OKESA- pinning techniques of the scarf type; OSHIHO – pinning techniques of the four corner hold; SRESP – respiratory chocking; SVASC – vascular chocking. \* Significant difference between males and females, <sup>#</sup> significant difference between group 1<sup>st</sup> and group 3<sup>rd</sup>.

**Table 3.** Number of penalties imposed during fights of men and women according to weight categories.

Instances of penalties and codes	Total	Males	Females	Male groups/weight categories				Female groups/weight categories			
				Group 1	Group 2	Group 3	Group 4	Group 1	Group 2	Group 3	Group 4
P29 Non-combativity	364	238	126 <sup>*</sup>	21	95	70	52 <sup>#,***,##</sup>	20	59	31	16
Other penalties	227	126	101	16	68	32	10	15	55	17	14
Total	591	364	227	37	163	102	62	35	114	48	30

\* Significant difference males from females, <sup>#</sup> Significant difference between group 1<sup>st</sup> and group 4<sup>th</sup>, \*\*\* significant difference between group 2<sup>nd</sup> and group 4<sup>th</sup>, ## significant difference between group 3<sup>rd</sup> and group 4<sup>th</sup>.

DISCUSSION

Hierarchy of throw techniques scored by male and female judo Olympians

Judo athletes preferred PLmaxA techniques. They performed these techniques more often than PLmidA, particularly PLminA techniques. From the biomechanical point of view, the force with the same magnitude and direction that acts on the greater lever causes greater effect (moment of force). The frequency of performing the above techniques depended neither on gender nor on weight category. This status reflects the principle that is used in technical and tactical preparation of judokas i.e. „Maximum-Efficiency with Minimum Effort” [5]. The underlying idea of judo declares the possibility of winning with opponents with greater physical strength. According to this principle, technical excellence means using the strength and inertia of the opponent against them [2]. In general, at equal resistance, when the arm of the lever used in a lever technique increases the applied

force decreases. This means that lever techniques of maximum arm are energetically most effective among lever techniques group. But more subtle information can be derived from this analysis on fighting rhythm.

In general, throwing techniques are connected with shifting velocity of Athletes couple system during the fight. In fact, using whatever lever techniques *tori* (attacker) needs for a while to stop himself to properly apply the technique. In the last Olympic Games, the rhythm of a fight was relatively quiet, also caused by the high non-combativity (65.4% for males and 55.5% for females). Coordinative and strength athletes’ capabilities were also increased thanks to the increasingly advanced training methodologies. This happens because the lever techniques are more complex (Figure 1, in Annex) as the whole movement, and they need higher coordination of the body and kinetic chains [GAI + (SSAI + ISAI) + Lever + Kake], for example *seoi-nage* (Figure 2) compared to couple techniques or [GAI + Couple + Kake] *uchi-mata* (Figure 3), but they are also





**Figure 2.** Seoi-nage is typical technique of Physical lever applied with variable arm [8]. (David Finch with permission).

more energy-consuming, as already demonstrated in many specific papers [10,14–17].

Men used the PLvA technique more often than women. The lower frequency of the effective PLvA techniques used by women was probably caused by lower upper body strength reflected in bench press and rowing tests [18]. Another group of judokas was characterized by higher percentage of relative torque in knee extensors, with lower percentage of flexors and trunk extensors compared to untrained controls. Although judo contestants exhibit similar relative strength to untrained peers, many years of training cause that they demonstrate higher strength in the muscles that are active when pulling or lifting the opponent when performing throws. Antigravity muscles are able to develop particularly high force in these people: they play an essential role when throws are performed [19]. Individual body build characteristics and experience cause that strength profile in elite seniors was connected with the preferred techniques of performing throws (foot and leg techniques or hand techniques) [20]. An explanation of the difference observed in the frequency of physical variable arm level techniques (PLvA) between males and females can be provided with an example of a *seoi-nage* throw. When shoulder throws such as *seoi-nage* are performed, a compensation of body posture can be observed, connected with disproportions in the status of force development. With knee extensors weaker than hip extensors, smaller knee bend is naturally observed. Lifting opponents will occur with unfavorable position of inclination forward (longer lever arm for the acting force). This situation is typical of weak antigravity muscles in lower extremities, both knee and hip extensors. If an athlete's knee extensors are weaker than those in hips, this state can be compensated by higher hip bend angle, without the necessity to incline the body trunk [19]. The relatively high flexibility in female kinetic chains often compensate for the weaker knee extensors with the helping application



**Figure 3.** Uchi-mata is typical technique of Couple forces applied by trunk and leg [8] (David Finch with permission).

of *makikomi* supplementary movement in PLvA application, however the weaker arm strength and, in general, different hip and gluteus dimensions make it very difficult to use these techniques fast and explosively.

#### Differences in throw techniques scored between extra lightweight and heavy weight male athletes

Significant differences between men from extra lightweight and heavy weight categories were found in PLmaxA, PLvA and CTL techniques. In addition, an increase in contingency coefficient strength was observed between 1<sup>st</sup> and next consecutive weight categories, i.e. 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup>. The proportionality of stature that changes with weight category is likely to have a particular importance. Heavyweights are usually proportionally shorter than lightweights, i.e. they are less ectomorphic than lightweights [21]. The body proportionality of an athlete should be related to his/her techniques preferred [22]. It is essential for judo that a compromise between keeping optimal body weight and composition and both physiological and motor efficiency is obtained [23]. Many results obtained for fat percentage in judokas were evaluated using different equations. However, the research carried out by the same authors and using the same methods [24] demonstrated increased adiposity in judokas from heavier weight categories. Those results corroborated findings of Callister et al. [25], who found moderate correlations between body mass and percent fat.

Relative dimensions of trunk and the differences in body mass and relative arms' strength are related to the significant differences between men from extra lightweight and heavy weight categories application of PLmaxA, PLvA and CTL techniques. In general, the heavyweights like to apply couple techniques that are simpler and energy-efficient.

As mentioned above, lever techniques are more complex as the whole movement, but they also need higher

coordination of the body and kinetic chains [GAI + (SSAI + ISAI) + Lever + Kake] compared to the couple techniques. [GAI + Couple + Kake]. In terms of the fight, this means that heavyweights, who prefer quiet pace during a contest, apply these relatively simpler techniques with high velocity to shorten the distance and fast application of couple. The bigger trunk dimensions support the CTL use because it promotes essential mechanics of this kind of techniques (to move the heavy adversary's body around his center of mass). On the contrary, they obviously have more difficulty in applying both PLmaxA and PLvA than competitors from extra lightweight categories, because, in general, the coordinative capabilities are lesser than in the lighter categories, but also because the essential mechanics for the physical lever techniques is the result of a well-coordinated and well-interconnected action performed by both kinetic chains in different time sequences that aims to translate the adversary's centre of mass in space [10].

First, there is a superior-chain open space that involves the body as part of the opponent's grip; secondly, there is the general action (reducing the distance) that is pursued and harmonically followed up by the coordinated and connected work of both Inferior and Superior Action Invariants as achieved through the abdominal and trunk muscles. These techniques need more skill in harmonic chains-connected movements, than Couple techniques; in fact, such techniques are often ineffective because of a lack in harmony in one of the preceding movements halts the throw, essentially preventing any score. Obviously such harmonic-complex movements are easier for extra light weights than for heavy ones.

The body composition factor can interact with the preference for a particular technique performed by heavyweights or lightweights, as relative strength tends to be frequently lower in heavyweights than in lightweights. High resistance in training and competition during many years of sport-selection process is likely to cause changes in body build. A very low difference of sexual dimorphism index was observed for height-weight ratio, ectomorphy, fat free mass percentage and calf girth. Average index in untrained subjects was higher than in judokas [26]. More often for female athletes, application of throws is Innovative or Classic, very few Chaotic Forms [10] are seen in women competition, but the percentage of Innovative variations is higher due to their body's flexibility. Connection *tachi-waza* → *ne-waza*, for *koshi-waza* is very often linked to the application of *makikomi* variation of throwing techniques. Normally, in women competitions, grip fight is less strength-based, while the attack velocity is not as explosive as in men competition. It is interesting to note that the poor presence of Chaotic Form of techniques in women games is directly

connected to the natural and relative lack of strength both in hands and legs of female athlete's body structure. Therefore, women's judo remains more connected to classic Kodokan Judo as for grips preference and the form of throwing techniques applied (Classic or Innovative) [10].

Relatively more vascular chokes instances were observed in women compared to men. This is likely to be connected to the unified training methods that aim to increase arms strength both in male and female athletes for grip goals. It is common knowledge that the muscular force generated in arms by women could increase with strength training to the level of 85–90% of the values recorded in men with similar weight, although this information has not been validated by any scientific studies. Recent studies, however, [27] found that female elite athletes (involving well-trained judo athletes) had lower hand grip strength than the untrained male subjects.

Furthermore, the change in arm strength is very often not connected to the similar increase in other muscle groups like neck, which in female athletes is probably weaker, with less developed sterno-cleido-mastoid muscles that protect carotid arteries in women and man. This problem needs a biomechanical research in a future.

Unexpectedly, the frequency of penalties imposed for non-combativity was significantly higher in the group of men than in women. It can be associated with the differences in psychological preparation rather than physical one [28,29].

## CONCLUSIONS

Because the techniques preferred during fighting depends on a gender and weight category of judokas, the relationships found in this study, which can be justified by the biomechanics of throws, should be taken into consideration in technical and tactical coaching of the contestants. A method used in this study can be recommended for future research concerning coaching tendencies.

## Practical applications


Statistical relationships concerning the choice of the fighting technique depending on gender and weight category were justified with biomechanics of the throws performed. Normally, it is well known that couple techniques are energetically more convenient compared to lever techniques. Body build should be considered when choosing the fighting technique, particularly when the opponent is higher or shorter or they use an opposite left or right grip *kenka-yotsu*. The quality of actions performed by the contestants should be monitored and analysed during competitions and training in order to optimally select the means of physical preparation and

stimulate technical and tactical preparation in terms of counterattack techniques or combined techniques. In order to achieve this, it is essential to focus on individual training of a particular contestant.

Throughout the years, female judo (which was biomechanically more Kodokan classic) have approached the

men's style. The observation of the London Olympic Games indicates a very unified approach to the training methodologies among male and female athletes in the world, that highlight the equivalent PLmaxA and CAL percentage in a fight, in spite of the natural differences in arm strength between male and female athletes.

ANNEX



**Figure 1.** Summary of the Kuzushi Tsukuri Action Invariants connected to Kake phase and Classic or Innovative and New (or Chaotic) Form of throwing techniques [10]. The figure is based on the most recent Kodokan classification [11] and innovative techniques names [12] and others.

**Table 4A.** Technique based on a couple of forces [8].

"Couple of Forces"- type Throwing Techniques	Couple applied by:	Arms	<i>Kuchiki-daoshi, kibisu-gaeshi, kakato-gaeshi, te-guruma, uchi-mata-sukashi</i>	All Innovative Variations of Throws and very few Chaotic Forms of Throws
		Arm/s and leg	<i>De-ashi-barai, o-uchi-gari, okuri-ashi- barai, ko-uchi-gake, ko-uchi-barai, ko- soto-gake, o-uchi-barai, harai-tsuri-komi-ashi, tsubame-gaeshi, yoko-gake, ko-uchi-gari, o-soto-gake, ko-soto-gari, o-uchi-gake, o-uchi-gaeshi (1)</i>	
		Trunk and legs	<i>O-soto-gari, o-tsubushi, o-soto-guruma, o-soto-otoshi, uchi-mata, ko-uchi-sutemi, okuri-komi-uchi-mata, harai-makikomi, harai-goshi, ushiro-uchi-mata, ushiro-hiza-ura-nage, hane-goshi, gyaku-uchi-mata, hane-makikomi, daki- ko-soto-gake, yama-arashi (Khabarelli- type throw), uchi-mata-gaeshi, hane-goshi gaeshi, harai-goshi-gaeshi, uchi-mata-makikomi, hane-makikomi</i>	
		Trunk and arms	<i>Morote-gari</i>	
		Legs	<i>Kani-basami</i>	

The table is based on the most recent Kodokan classification [11] and innovative techniques names [12]

**Table 4B.** Technique based on a physical lever [8].

Physical Lever- type Throwing Techniques Lever applied by:	Minimum Arm Lever (fulcrum under uke's waist)	<i>O-guruma, ura-nage, kata-guruma, ganseki-otoshi, tama-guruma, uchi-makikomi, binta-guruma, obi-otoshi, soto-makikomi, tawara-gaeshi, makikomi, kata-sode-ashi-tsuru, sukui-nage, daki-sutemi, ushiro-goshi, utsuri-goshi</i>	All Innovative Variation and New (Chaotic) Forms
	Medium Arm Lever (fulcrum under uke's knees)	<i>Hiza-guruma, ashi-guruma, hiza- soto-muso, soto-kibisu-gaeshi</i>	
	Maximum-Arm Lever (fulcrum under uke's malleolus)	<i>Uki-otoshi, yoko-guruma, yoko-otoshi, yoko-wakare, sumi-otoshi, seoi-otoshi, suwari-otoshi, hiza-seoi, no-waki, o-uchi-gaeshi (2) waki-otoshi, obi-seoi, tani-otoshi, suso-seoi, tai-otoshi, suwari-seoi, dai-sharin, hiza-tai-otoshi, hikkomi-gaeshi, tomoe-nage, sumi-gaeshi, ryo-ashi-tomoe, yoko-kata-guruma, yoko-tomoe, uki-waza, sasae-tsuru-komi-ashi, uke-nage</i>	
	Variable Arm (variable fulcrum from uke's waist to his knees)	<i>Tsuru-komi-goshi, kubi-nage, o-goshi, sasae-tsuru-komi-goshi, koshi-guruma, ko-tsuru-komi-goshi, o-tsuru-komi-goshi, sode-tsuru-komi-goshi, seoi-nage, eri-seoi-nage, uki-goshi, morote-seoi-nage</i>	

The table is based on the most recent Kodokan classification [11] and innovative techniques names [12].

REFERENCES:

1. IJF Extraordinary Congress, Refereeing rules alterations, Bangkok, 26th October 2008, Thailand. Available from URL: [http://www.intjudo.eu/?Menu=News&Action=Full&type\\_id=1&sid=146&lang\\_id=2&mid=4](http://www.intjudo.eu/?Menu=News&Action=Full&type_id=1&sid=146&lang_id=2&mid=4)

2. Sterkowicz S, Kiejda I, Błach W: Charakterystyka sposobów walki judo w Igrzyskach Olimpijskich 1988–1996 [Fighting ways in judo during Olympic Games 1988–1996]. *Trening*, 1997; 1: 27–41 [in Polish]

3. Sterkowicz S, Franchini E: Techniques used by judoists during the world and Olympic tournaments 1995–1999. *Human Movement*, 2000; 2(2): 23: 33

4. Sterkowicz S: Differences in the specific movement activity in men and women practicing judo. *J Hum Kinet*, 1999; 1: 99–113

5. Illustrated Kodokan Judo. In: Kodansha, editor. Tokyo, Japan, 1964

6. Sterkowicz S, Lech G, Almansba R: The course of fight and the level of sports achievements in judo. *Arch Budo*, 2007; 3: 72–81

7. Sacripanti A: Biomechanical classification of judo throwing techniques (Nage Waza). 5th International Symposium of Biomechanics in Sport, Greece: Athens; 1987; 181–94. Available from URL: <http://cdsweb.cern.ch/record/1111755>

8. Sacripanti A: Advances in Judo Biomechanics Research “Modern Evolution on Ancient Roots”, VDM Verlag Dr. Muller, Sarbrucken, 2010

9. Official results book JUDO 28 July – 3 August, London Organising Committee of the Olympic Games and Paralympic Games Limited, London, 2012; 3007–138

10. Sacripanti A: A Biomechanical Reassessment of the Scientific Foundations of Jigorō Kanō’s Kōdōkan Jūdō, Cornell University Library. Available from URL: <http://arxiv.org/abs/1206.1135>

11. Daigo T: Kodokan Judo Throwing Techniques, Kodansha International Ltd., USA, 2005

12. Inman R: Shin kokusai shiai waza. University of Bath, 2007

13. Stupnicki R: Biometria/Biometry. Margos, Warszawa, 2005

14. Franchini E, de Moraes C, Bertuzzi R et al: Energy expenditure in different judo throwing techniques. Proceedings of first Joint International Pre-Olympic Conference of Sports Science and Sports Engineering: Nanjing, China, August 4–7, 2008, 2: 55–60

15. De Blasis T, Rossetti E, Andreatta G et al: Infrared Thermography- calorimetric Quantitation of Energy Expenditure in Biomechanically Different Types of Jūdō Throwing Techniques. 3rd European Science of Judo Poster Exhibition, European Judo Union, Chelyabinsk, 26 April 2012: 4–5

16. Peixoto R, Monteiro L: Structural analysis and energetic comparison between two throwing techniques uchi-mata vs ippon-seoi-nage. 3rd European Science of Judo Poster Exhibition, European Judo Union, Chelyabinsk 26 April 2012: 28–29

17. Sacripanti A: Kumi Kata “from dojo to high level competition”. [book in press]

18. Monteiro L, García García JM, Carratala V: The balance and the ratio of pulling-reaping-pushing in elite judokas. In: Proceedings (editors Figueiredo AA, Gutiérrez-García C.) Scientific Congress on Martial Arts and Combat Sports. Associação para o Desenvolvimento e Investigação de Viseu, Instituto Politécnico de Viseu Escola Superior de Educação de Viseu, Viseu – Portugal 13<sup>th</sup>–15<sup>th</sup> May 2011; 77–81

19. Sterkowicz S, Lech G, Chwała W et al: Muscle strength in young judo contestants vs. untrained subjects. *Arch Budo*, 2011; 7(3): 179–84

20. Dąbrowska A, Sikorski W, Wit A: Dyspozycje siłowe zawodnika judo a skuteczność ich techniki [Power and technique effectiveness of judo athletes]. *Sport Wyczynowy*, 1986; 3–4: 35–42 [in Polish]

21. Claessens A, Beunen G, Simons JM et al: Body structure, somatotype, and motor fitness of top-class Belgian judoists. In: The 1984 Olympic Scientific Congress Proceedings: Perspectives in Kinanthropometry, (ed.) Day JAP, Human Kinetics Publishers, Champaign, Illinois, 1984; 155–63

22. Detanico D, Santos SG: A Relação Entre a Proporcionalidade Corporal do Judoca e sua Técnica de Preferência (Tokui-Waza). *R. bras. Ci e Mov*. 2007; 15(3): 15–24 [in Portuguese]

23. Almansba R, Sterkowicz S, Belkacem R et al: Anthropometrical and physiological profiles of the Algerian Olympic judoists. *Arch Budo*, 2010; 6(4): 185–93

24. Iida E, Nakajima T, Wakayama H, Matsumoto D: Rating scales of fundamental physical fitness for college judoists: composition and application. In: National Judo Conference. International Research Symposium Annals. United States Olympic Training Center, Colorado Springs, 1998; 13

25. Callister R, Callister RJ, Staron RS et al: Physiological characteristics of elite Judo athletes. *Int J Sports Med*, 1991; 12(2): 196–203

26. Sterkowicz-Przybycień K, Almansba R: Sexual Dimorphism of Anthropometrical Measurements in Judoists vs. Untrained Subject. *Science & Sports*, 2011; 26: 316–23

27. Leyk D, Gorges W, Ridder D et al: Hand-grip strength of young men, women and highly trained female athletes. *Eur J Appl Physiol*, 2007; 99: 415–21

28. Blumenstein B, Lidor R, Tenenbaum G: Periodization and planning of psychological preparation in elite combat sport programs: the case of judo. *IJSEP*, 2005; 3: 7–25

29. Sterkowicz S, García García JM, Suay, Lerma F: The importance of judo trainers’ professional activities. *Arch Budo*, 2007; 3: 57–61

Cite this article as: Sterkowicz S, Sacripanti A, Sterkowicz-Przybycień K: Techniques frequently used during London Olympic judo tournaments: A biomechanical approach. *Arch Budo*, 2013; 1: 51–58.