

# Counter movement jump performance between older adults with and without regular tai chi exercise training

Jung-Chun Chi<sup>1BD</sup>, Chen-Fu Huang<sup>1ADE</sup>, Thomas W. Kernozek<sup>2CD</sup>, Yao-Yi Hsieh<sup>1BD</sup>

## Authors' Contribution:

- A Study Design
- B Data Collection
- C Statistical Analysis
- D Manuscript Preparation
- E Funds Collection

<sup>1</sup>Department of Physical Education, National Taiwan Normal University, Taipei, Taiwan

<sup>2</sup>Department of Health Professions, Physical Therapy Program, University of Wisconsin-La Crosse, La Crosse, Wisconsin

**Source of support:** "Aim for the Top University Plan" of the National Taiwan Normal University and the Ministry of Education, Taiwan, R.O.C.

**Received:** 03 January 2013; **Accepted:** 07 November 2013; **Published online:** 29 November 2013

**ICID:** 1078408

## Abstract:

**Background & Study Aim:** Martial arts, especially tai chi and other systems of health exercises can significantly stimulate all dimensions of the positive health. The aim of the study was the effect of tai chi exercise on maximal counter movement jump (CMJ) in older adults.

**Material & Methods:** Tai chi group (n=17; age: 72.1 ± 4.3 yrs) and healthy older adults (n=17; age: 73.5 ± 5.5 yrs) were recruited in this study. Jump height, vertical velocity of the body center of mass (COM), and peak power output were analyzed by using a force plate (1000Hz) during the performance of a maximal effort CMJ.

**Results:** Tai chi subjects had greater performance than healthy elderly group in jump height (14.35±4.07cm vs. 11.18±3.41cm), peak power output (29.72 ± 3.82 vs. 25.52 ± 4.04 watts/BW), vertical velocity of the body COM at takeoff (V<sub>take-off</sub>) (1.68 ± 0.20 vs. 1.49 ± 0.22 m/sec).

**Conclusions:** Older adults participated in tai chi exercise training had better performance in a maximal effort CMJ. These enhancements may decrease limitations of daily functional performance and reduce fall risk.

**Key words:** center of mass • fall prevention • kinematics • lower extremity • martial arts • muscle power

**Author's address:** Chen-Fu Huang, Department of Physical Education, National Taiwan Normal University, No. 88, Sec. 4, Tingshou Rd., Wenshan Dist., Taipei City 116, Taiwan (R.O.C.); e-mail: t08001@ntnu.edu.tw

**Positive health** – a concept of health related to the quality of life and to the capability possessed by an individual. This term relates more to the development, than to the simple coping skills. In the physiological context it may be perceived as a state which is characterized by: A) the absence of the disease; b) low level of the severity of risk factors of the civilization diseases; c) an adequate capacity of adaptive mechanisms responsible for the control of the external environment, the physical effort in particular [31].

## INTRODUCTION

Age-related decline in muscle function, including muscle strength and muscle power, increases the difficulty in the performance of daily functional activities in older adults. The rate of decline in muscle power after the age of 65 years has been reported to be higher than the overall reduction in muscular strength [1]. As a result, this overall decrease in muscle power may have a greater impact on functional performance of daily activities than the reduction in muscle strength [2-4]. Reduced muscle power has been cited as one of the primary factors that may impair mobility and maintain

independence of older adults [5-7]. Results of previous studies have also reported increased frailty, a reduction in postural control and increased rate of falls in adults of increased age [3,8,9].

Regular exercise is essential for older adults in order to maintain muscle function. Tai-chi exercise, a traditional Chinese martial art, is a very popular and accepted exercise modality in both eastern and western countries. Many studies have reported that engaging in tai chi exercise may enhance strength [10-13], flexibility [14], and balance [15-17]. Tai chi exercise

programs for older adults have been often designed to combine strength, flexibility and balance training [8,18]. Furthermore, the moderate intensity of tai chi exercise [19] is possible for older adults. In our opinion, tai chi exercise may be suitable for older adults to engage in terms of physical enhancement, training intensity, and the forms of body movements required. In addition, tai chi exercise can be performed just about anywhere and with little specialized equipment.

The countermovement jump (CMJ), a test of physical performance, is a dynamic weight bearing movement skill that utilizes multiple muscles and joints of the lower extremities. This assessment could be used to estimate muscle strength and power performance in older adults [20,21]. Both components of muscle power, muscle force and contraction velocity, can be observed during a maximal CMJ [22,23]. Investigations [22,23] have suggested that the CMJ, can detect significant functional limitations in muscle strength and power of older adults during this coordinated motor skill. Muscle power production through pre-stretch enhancement during this weight-bearing skill in older adults may be highly relevant to their functional performance capabilities.

Although there are many studies that have examined tai chi exercise's influence on the specific functional tasks such as gait [24] and balance [25,26] in older adults, to our knowledge, there appears to be no evaluation of maximal performance effort using a CMJ in older adults that regularly perform tai chi exercise. Previous study suggested that it may be of interest to examine CMJ performance in older adults that have regularly performed regular tai chi exercise [23]. Many studies have reported that muscle strength of lower limbs could be enhanced through tai chi training [10-13,27,28]. However, these studies have largely examined isometric or isokinetic strength of single lower extremity joints [10-13]. These strength assessments do not address muscle power output during a dynamic, multijoint, weightbearing task such as required in the CMJ.

The aim of the study was the effect of tai chi exercise on the maximal counter movement jump (CMJ) performance in older adults. We compared between regular tai chi elderly performers and healthy elderly adults during maximal CMJ to determine if there were changes in muscular strength and power of the lower extremities with tai chi exercise. We hypothesized that long-term tai chi exercise group would have a greater CMJ performance than the healthy older adult group.

## MATERIAL AND METHODS

Seventeen older adults who were regular tai chi exercise performers (TC older adult group: included 10 males and 7 females; average age:  $72.12 \pm 4.31$  yrs; average height:  $159 \pm 6.74$  cm; average weight:  $57.76 \pm 6.89$  kg) and 17 older adults without regular and systematic physical training (healthy older adult group: included 11 males and 6 females; average age:  $73.5 \pm 5.51$  yrs; average height:  $164.38 \pm 8.77$  cm; average weight:  $61.16 \pm 9.93$  kg) were recruited. All subjects of TC older adult group regularly performed traditional *Yang Tai Chi* exercise (108 forms) at least 5 times per week for more than two years. TC exercise sessions consisted of approximately 10-min TC warm-up, 25-min of main TC exercise, and 5-min of cool-down. Participants who were currently suffering from motor or neuropsychological disorders were not included in this investigation. All subjects had adequate warm-up prior to testing in the laboratory. This research was approved by the local institutional review board and all volunteers signed a written consent form prior to testing.

A Kistler 9281 force plate (Kistler Instrumente AG, Winterthur, Switzerland, 90 cm in length, 60 cm in width) interfaced with a computer was used to record the ground reaction forces (GRF) at 1000Hz while the subjects performed 3 maximal effort CMJs. Each trial was recorded from the beginning of the movement until contact with the force plate after the flight phase of the jump.

Subjects were asked to place and keep their hands at their hips while performing the maximal effort CMJ. Each subject was instructed to start in an upright position, rapidly squat down to about 90 degrees of knee flexion and immediately perform a maximal jump into the air (Figure 1). They were asked to land back on the force plate during all performance trials. The downward depth and speed in which all subjects performed the CMJ was self-selected by the performer. For safety, an experienced research assistant stood alongside each subject while performing the test. One of three trials with the greatest time in the air was selected for further analysis. The vertical ground reaction force (VGRF) from the force platform was used to calculate the vertical velocity of the body center of mass (COM) according to the following equation.

$$Vc(t) = \frac{1}{m} \int_{t_0}^t [F(t) - mg] dt$$

Where  $Vc$  is the vertical velocity of COM, F is the vertical force measured by the platform, m is the body mass of the subject and g is the acceleration due to

gravity ( $9.81 \text{ m s}^{-2}$ ),  $t_0$  is the beginning of the CMJ, the instant that the VGRF was less than 10 N below body weight and decreased during the countermovement phase of the CMJ. Power of the lower extremities was continuously calculated during the CMJ as the instantaneous product of VGRF and  $V_c$  [23]. The position of COM was obtained by integrating instantaneous velocity during the CMJ.


The CMJ was divided into two phases: the countermovement phase and the jump phase. The countermovement phase was defined as the time interval from starting position to the point associated with the lowest position of COM. The jump phase was defined as the time interval from the lowest position of COM to the take-off from the force platform (Figure 1). Consequently, the duration of countermovement and jump phase was calculated by the time interval based on these two events, and the displacement of COM during phase was determined by change in vertical position of COM. The VGRF and power were normalized by each subjects' body weight. Based on the vertical velocity at the instant of take-off (i.e. take-off vertical velocity) and projectile motion principles, we calculated the jump height.

Independent t-tests were used to examine differences between the two groups (TC and non-TC

participants) for each dependent variable. Statistical analyses were performed using SPSS 12.0 statistical software (SPSS Inc., Chicago, IL, USA). Alpha was set to .05 for all statistical tests.

## RESULTS

Table 1 depicts the data for each group during the performance of the maximal CMJ. At the lowest position of COM during the countermovement phase, TC group exhibited 9.4% greater vertical ground reaction force (VGRF) than healthy older adult group, while the duration and displacement of COM during this phase were not different ( $p > .05$ ). During jump phase, the two groups also exhibited no difference in duration and displacement of COM ( $p > .05$ ). Compared to the healthy older adult group, TC group exhibited a greater jump height, peak vertical velocity ( $V_{\text{peak}}$ ), take-off vertical velocity ( $V_{\text{take-off}}$ ), peak VGRF, and peak power ( $p < .05$ ). Since peak power is obtained the greatest production of vertical force and the corresponding instantaneous vertical velocity, we showed that both components of peak power, VGRF and vertical velocity, were greater in TC exercise group. Finally, compared to the healthy older adult group, TC group exhibited a greater ratio of  $V_{\text{take-off}}/V_{\text{peak}}$  ( $p < .05$ ).



**Figure 1.** The CMJ was divided into two phases: the countermovement phase (from a to b) and jumping phase (from b to c). The a, b and c are the instant of beginning, the lowest position of COM, and take-off during CMJ, respectively.

**Table 1.** Variables of the tai chi (TC) training group and the healthy elder group during the maximal CMJ

Variable	TC elderly (n = 17)		healthy elderly (n = 17)		p
	M	SD	M	SD	
<b>Counter-movement phase</b>					
Duration (ms)	602.80	67.30	669.60	230.10	0.34
COM displacement (cm)	21.73	7.97	22.81	7.13	0.68
Peak vertical force (N/BW)	17.15	2.35	15.54	2.12	0.04*
<b>Jump phase</b>					
Duration (ms)	295.80	74.30	320.00	55.64	0.29
COM displacement (cm)	32.59	7.98	32.92	7.24	0.90
Peak vertical force (N/BW)	20.12	2.02	18.19	1.53	0.00*
Peak vertical velocity (m/s)	1.82	0.16	1.69	0.19	0.05*
Instantaneous velocity <sub>take-off</sub> (m/s)	1.68	0.20	1.49	0.22	0.02*
Peak power (W/BW)	29.72	3.82	25.52	4.04	0.00*
Velocity <sub>peak power</sub> (ms <sup>-1</sup> )	1.64	0.14	1.53	0.17	0.05*
Force <sub>peak power</sub> (N/BW)	19.59	2.02	16.53	1.31	0.00*
V <sub>take-off</sub> /V <sub>peak</sub> (%)	92.36	3.05	88.15	4.93	0.01*
Jump height (cm)	14.35	4.07	11.18	3.41	0.02*


\**p* < 0.05

## DISCUSSION

These findings support our hypothesis that the TC exercise group would have greater performance during a maximal CMJ, including jump height, the VGRF at the lowest position of COM during the counter-movement phase and peak VGRF, peak power, force and velocity components of peak power,  $V_{\text{peak}}$ ,  $V_{\text{take-off}}$ , ratio of  $V_{\text{take-off}}/V_{\text{peak}}$  during jumping phase.

The ground reaction force variables obtained represent the ability of neuromuscular system to develop strength and power in the lower extremities during a maximal effort CMJ [21]. In the present study, TC older adults had greater peak VGRF in the CMJ during the counter-movement and jumping phases compared to the healthy older adults who did not regularly participate in TC exercise training. These results reveal that older adults that regularly participate in TC exercise appear to have a better neuromuscular capacity to produce muscle strength and power during a maximal CMJ. It is well known that muscle strength and power tends to decrease with ageing. Skelton et al. [1] reported a decline in muscle strength of 1.2~2% per year after the age of 65 years. In our study, those who performed regular and long-term TC exercise may have enhanced or maintained muscle strength and power. This may reduce some of the age related limitations of associated with functional activities.

De Vito et al. [29] estimated the maximal power of sedentary women aged 50-75 years through two types of vertical jumps: squat jumps (SJ) and counter-movement jumps (CMJ). They reported that maximal muscle power decreased significantly with aging. In their study, another main finding was that the instantaneous vertical velocity associated with the maximal power was negatively and significantly related to these older adults while the magnitude of VGRF was not related to age. In our investigation, the TC exercise group developed significantly greater peak power during a maximal effort CMJ than the healthy older adult group (Figure 2a and Figure 2d). This increase in peak power may due to the effect of regular long-term traditional Yang's Tai Chi exercise. An increase in peak power in older adults has also been observed after both a 36-week multi-component training [22] and a 12 week of low intensity conditioning exercise programs [30] in other investigations. In a further examination of the VGRF and instantaneous velocity components at the instant of peak power, there appears to be a significant increase in the instantaneous velocity with no change in VGRF at the instant of peak power in either of these investigations (about 4.1 % and 13% change, respectively). The most novel part of our investigation was that the TC elderly group when compared to



**Figure 2.** Left column (a, b, c) and right column (d, e, f) are the typical power (watt/body weight), instantaneous velocity of COM (m/s), and vertical ground reaction force (N/BW) of a Tai Chi and healthy older adult, respectively. The CMJ was normalized as 100% from the start to the take-off. The instant of peak power was denoted at the vertical line shown at PP.

the healthy older adult group had a greater VGRF and instantaneous velocity at the instant of peak power (Table 1). This may have been due to the regular participation (greater than 2 years) in Tai Chi exercise of the TC participants in our investigation.

The velocity variables,  $V_{\text{peak-power}}$ ,  $V_{\text{peak}}$ , and  $V_{\text{take-off}}$ , were greater in TC exercise group compared to the healthy older adults in this study. These results demonstrated that the movement velocity was enhanced in the TC exercise group. These changes influenced the power output and resulting vertical jump performance. The TC exercise group showed a greater ratio of  $V_{\text{take-off}}/V_{\text{peak}}$  compared to the healthy older adult group.  $V_{\text{peak}}$  occurred during the last portion of CMJ prior to take-off (Figure 2b and Figure 2e). The decreases in instantaneous velocity from the  $V_{\text{peak}}$  to  $V_{\text{take-off}}$  time interval may represent the group's ability of accelerate their center of mass through their lower extremities. A higher ratio of  $V_{\text{take-off}}/V_{\text{peak}}$  in the TC exercise group may be a reflection of better muscle capacity prior to take-off. Caserotti et al. [23] reported that the ankle plantar flexors may be responsible for these findings. Previous investigations have reported increases in ankle muscular

strength in tai chi training groups [27,28]. However, we are not sure if the higher ratio of  $V_{\text{take-off}}$  and  $V_{\text{peak}}$  observed in the TC exercise group in our study could be attributed to stronger plantar flexors. Therefore, further investigation using kinematics and EMG measures may be useful in further examining this hypothesis.

Besides jump height, the TC exercise group showed many performance increases in our measurements. The specific movements, such as one-leg deep squat, two-leg deep squat and jumping, are required in 108-forms of traditional Yang's Tai Chi exercise. Many of these movement forms appear to provide an exercise training stimulus for many muscle groups in a weight bearing situation that improve the performance of a CMJ.

## CONCLUSIONS

The older adults who participated in regular TC exercise exhibited greater performance (including greater peak VGRF, peak power, peak vertical velocity, take-off velocity, and jump height) during a maximal CMJ compared to a group of healthy older



adults. Both the VGRF and instantaneous velocity at time of peak power was enhanced in our TC exercise group. The TC exercise group exhibited a better capacity to maintain a higher peak vertical velocity of COM prior to take-off. These performance enhancements may influence the functional performance of elderly individuals.

## ACKNOWLEDGEMENTS

This work is particularly supported by “Aim for the Top University Plan” of the National Taiwan Normal University and the Ministry of Education, Taiwan, R.O.C.

## REFERENCES

- Skelton DA, Greig CA, Davies JM et al. Strength, power, and related functional ability of healthy people aged 65–89 years. *Age Ageing* 1994; 23: 371–377
- Bassy EJ. Measurement of muscle strength and power. *Muscle Nerve* 1997; 20(5): 44–46
- Bassy EJ, Fiatarone MA, O'Neill EF et al. Leg extension power and function performance in very old men and women. *Clin Sci* 1992; 82: 321–327
- Rantanen T, Era P, Heileleinen E. Maximal isometric strength and mobility among 75-year-old men and women. *Age Ageing* 1994; 23: 132–137
- Foldvari M, Clark M, Laviolette LC et al. Association of muscle power with functional status in community-dwelling elderly women. *J Gerontol A-Biol* 2000; 55(4): 192–199
- Rantanen T, Avela J. Leg extension power and walking speed in very old people living independently. *J Gerontology* 1997; 52A(4): 225–231
- Suzuki T, Bean JF, Fielding RA. Muscle power of the ankle flexors predicts functional performance in community dwelling older women. *J Am Geriatr Soc* 2001; 49: 1161–1167
- Fielding RA, LeBrasseur NK, Cuoco A et al. High-velocity resistance training increases skeletal muscle peak power in older women. *J Am Geriatr Soc* 2002; 50(4): 655–662
- Skelton DA, Kennedy J, Rutherford OM. Explosive power and asymmetry in leg muscle function in frequent fallers and nonfallers aged over 65. *Age ageing* 2002; 31(2): 119–125
- Christou EA, Yang Y, Rosengren KS. Taiji training improves knee extensor strength and force control in older adults. *J Gerontol A Biol Sci Med Sci* 2003; 58(8): 763–766
- Lan C, Lai JS, Chen SY et al. 12-month Tai Chi training in the elderly: Its effect on health fitness. *Med Sci Sports Exerc*, 1998 30(3): 345–351
- Jacobson BH, Chen HC, Cashel C et al. The effect of Tai Chi Chuan training on balance, kinesthetic sense, and strength. *Percept Mot Skills* 1997; 84(1): 27–33
- Qin L, Choy W, Leung K et al. Beneficial effects of regular Tai Chi exercise on musculoskeletal system. *J Bone Miner Metab* 2005; 23: 186–190
- Jones AY, Dean E, Scudds RJ. Effectiveness of a community-based Tai Chi program and implications for public health initiatives. *Arch Phys Med Rehabil* 2005; 86(4): 619–625
- Lin YC, Wong AM, Chou SW et al. The effects of Tai Chi Chuan on postural stability in the elderly: preliminary report. *Chang Gung Med J* 2000; 23(4): 197–204
- Schaller KJ. Tai Chi Chih: an exercise option for older adults. *J Gerontol Nurs* 1996; 22(10): 12–17
- Wong AM, Lin YC, Chou SW et al. Coordination exercise and postural stability in elderly people: Effect of Tai Chi Chuan. *Arch Phys Med Rehabil* 2001; 82(5): 608–612
- Izquierdo M, Häkkinen K, Ibañez J et al. Effects of strength training on muscle power and serum hormones in middle-aged and older men. *J Appl Physiol* 2001; 90: 1497–1507
- Li JX, Hong Y, Chan KM. Tai Chi: physiological characteristics and beneficial effects on health. *Br J Sports Med* 2001; 35: 148–156
- Bosco C, Cardinale M, Tsarpela O et al. The influence of whole body vibration on jumping performance. *Biol Sport* 1998; 15: 157–164
- Liu Y, Peng CH, Wei SH et al. Active leg stiffness and energy stored in the muscles during maximal counter movement jump in the aged. *J Electromyogr Kines* 2006; 16(4): 342–351
- Caserotti P, Aagaard P, Puggaard L. Changes in power and force generation during coupled eccentric-concentric versus concentric muscle contraction with training and aging. *Eur J Appl Physiol* 2008; 103(2): 151–61
- Caserotti P, Aagaard P, Simonsen EB et al. Contraction-specific differences in maximal muscle power during stretch-shortening cycle movements in elderly males and females. *Eur J Appl Physiol* 2001; 84(3): 206–212
- Hass CJ, Gregor RJ, Waddell DE et al. The influence of Tai Chi training on the center of pressure trajectory during gait initiation in older adults. *Arch Phys Med Rehabil* 2004; 85: 1593–1598
- Gatts SK, Woollacott MH. How Tai Chi improves balance: Biomechanics of recovery to a walking slip in impaired seniors. *Gait posture* 2007; 25: 205–214
- Li FH, Harmer P, Fisher J et al. Tai Chi: improving functional balance and predicting subsequent falls in older persons. *Med Sci Sports Exerc* 2004; 36(12): 2046–2052
- Li JX, Xu DQ, Hong Y. Changes in muscle strength, endurance, and reaction of the lower extremities with Tai Chi intervention. *J Biomech* 2009; 42: 967–971
- Xu DQ, Hong Y, Li JX. Tai Chi exercise and muscle strength and endurance in older people. *Med Sport Sci* 2008; 52: 20–29
- De VG, Bernardi M, Forte R et al. Determinants of maximal instantaneous muscle power in women aged 50–75 years. *Eur J Appl Physiol Occup Physiol* 1998; 78(1): 59–64
- De VG, Bernardi M, Forte R et al. Effects of a low-intensity conditioning programme on VO<sub>2</sub>max and maximal instantaneous peak power in elderly women. *Eur J Appl Physiol Occup Physiol* 1999; 80(3): 227–232
- Kuński H. Trening zdrowotny osób dorosłych. Poradnik lekarza i trenera. Agencja Wydawnicza Medsportpress. Warszawa 2002 [in Polish]

**Cite this article as:** Chi JC, Huang CF, Kernozek TW et al. Counter movement jump performance between older adults with and without regular tai chi exercise training *Arch Budo* 2013; 4: 249–254