

# Heart rate, a rating of perceived exertion and basic affective responses during different moments of a single *capoeira* progressive training session (CPTS)

## Authors' Contribution:

- ✍ A Study Design
- 📁 B Data Collection
- 📊 C Statistical Analysis
- 📄 D Manuscript Preparation
- 🏆 E Funds Collection

Sérgio R Moreira<sup>1,2,3ACDE</sup>, Alfredo A Teixeira-Araujo<sup>1,2BD</sup>, Mariana V F Camara<sup>3BD</sup>,  
Ferdinando O Carvalho<sup>1,2,3BE</sup>, Thaynã A Bezerra<sup>2B</sup>, Anastácio N Souza Filho<sup>2B</sup>,  
Karina E S Silva<sup>4D</sup>, Eduardo C Costa<sup>5D</sup>

<sup>1</sup> Graduate Program on Physical Education, Federal University of Vale do São Francisco – UNIVASF, Petrolina, Brazil

<sup>2</sup> Graduate Program on Health and Biological Sciences, Federal University of Vale do São Francisco – UNIVASF, Petrolina, Brazil

<sup>3</sup> Tutorial Education Program (PET-Educação Física) of Federal University of Vale do São Francisco – UNIVASF, Petrolina, Brazil

<sup>4</sup> Graduate Program on Physical Education of State University of Londrina – UEL, Londrina, Brazil

<sup>5</sup> Graduate Program on Physical Education of Federal University of Rio Grande do Norte – UFRN, Natal, Brazil

Received: 17 March 2017; Accepted: 28 April 2017; Published online: 27 December 2017

AoBID: 11530

## Abstract

### Background & Study Aim:

Little information is available on psychophysiological responses in martial arts which do not have the status of Olympic sport (such as judo, taekwondo) and yet are very popular. Lack of information about the impact of any effort on the human body means the lack of rational control over a given psychomotor human activity. The aim was the knowledge about heart rate (HR), rate perceived exertion (RPE) and basic affective responses during different moments of a single *capoeira* progressive training session (CPTS).

### Material & Methods:

Sixteen adults (10 male, 6 female) were evaluated to HR, rate perceived exertion (RPE 6/20), feeling scale (FS from 5+ to 5-) and perceived activation (PA six points scale). The CPTS lasted 60-min, being: 2-min/playing instruments; 2-min/singing; 8-min/warm-up; 8-min/*ginga* (the basic movement); 8-min/technical movements; 12-min/passive *capoeira*; 1-min/*capoeira* game. The remaining time was distributed in recovery during CPTS. A cycle ergometer test was performed starting at 25W/50W (female/male) with increments of 25W every 1min until exhaustion (50-RPM's) to anaerobic threshold (AT) and maximal power ( $P_{MAX}$ ) determination.

### Results:

When compared to resting the HR ( $77 \pm 13$  bpm) increased from warm-up ( $102 \pm 12$  bpm) until *capoeira* circle ( $135 \pm 25$  bpm), as well as differed from the total CPTS ( $117 \pm 13$  bpm). The total CPTS occurred at  $80 \pm 10\%$ HR-AT and  $65 \pm 7\%$ HR- $P_{MAX}$ . The RPE shows increase from resting ( $8 \pm 3$  points) to *ginga* ( $12 \pm 2$  points), technical movements ( $14 \pm 2$  points), *capoeira* circle ( $14 \pm 2$  points) and total CPTS ( $11 \pm 2$  points). The PA increased from resting ( $3 \pm 1$  points) to singing ( $4 \pm 1$  points), *ginga* ( $4 \pm 1$  points), technical movements ( $5 \pm 1$  points) and *capoeira* circle ( $5 \pm 1$  points).

### Conclusions:

A single CPTS increased the HR, RPE and PA responses, maintaining affective valence (FS) in the practitioners during the session. Besides, a CPTS may be an alternative physical activity modality to improve fitness and health of adults with physical activity intensity level (i.e., moderate-vigorous).

### Keywords:

emotions • exertion • *ginga* • internal load • perception • singing • workload

### Copyright:

© 2017 the Authors. Published by Archives of Budo Science of Martial Arts and Extreme Sports

<b>Conflict of interest:</b>	Authors have declared that no competing interest exists
<b>Ethical approval:</b>	The study was approved by the Research and Ethics Committee of the Federal University of Vale do São Francisco (protocol no 0001/200813 CEDEP)
<b>Provenance &amp; peer review:</b>	Not commissioned; externally peer reviewed
<b>Source of support:</b>	CAPES (Research Support Foundation of Brazil) to fund scholarships
<b>Author's address:</b>	Sérgio Rodrigues Moreira, Physical Education Department, Federal University of Vale do São Francisco, Av. José de Sá Maniçoba, S/N - Centro - CEP: 56.304-917 Petrolina, Brazil; e-mail: sergio.moreira@univasf.edu.br

**Ginga** – the basic movement of *capoeira* (the purpose of the *ginga* is not to offer the opponent a fixed target).

**Capoeira** – athletic sport composed of attack and defence systems created in colonial Brazil.

**Colonial Brazil** – comprises the period from 1500, with the arrival of the Portuguese, until 1815, when Brazil was elevated to a kingdom alongside Portugal.

**Basic Programmed Lesson** – practical method that allows the initiation of novices without any experience in *capoeira*.

**Atabaque and pandeiro** – characteristic rhythmic instruments of *capoeira* with the membrafonic design.

**Singing capoeira** – typical music of *capoeira*.

**Chronic** – adjective 1. used for describing a disease or condition that lasts for a long time. Compare **acute** 2. used for describing severe pain [31].

**Exertion** – noun physical activity [31].

**Perception** – noun the process of using the senses to acquire information about the surrounding environment or situation [31].

**Psychomotor** – adjective relating to bodily movement triggered by mental activity, especially voluntary muscle action [31].

## INTRODUCTION

*Capoeira* is a Brazilian martial art that has spread worldwide [1]. Currently, *capoeira* is present in more than 150 countries [2], and different social groups have carried out its practice over the five continents [3]. It was originally created in colonial Brazil and as a modality can be defined as an athletic sport characterised by attack and defence systems [4, 5]. The main body movement of *capoeira* is called “*ginga*”. Moreover, common actions of *capoeira* include dodge, unbalance, impact acrobatic movements [4], which are performed following *Angola*, *Benguela*, and *São Bento Grande* rhythms [6]. *Angola* and *Benguela capoeira* styles are more likely to rely on aerobic energy pathways, while *São Bento Grande* may alternate aerobic or anaerobic predominance, depending on the practitioner’s training status. Also, it is important to highlight that *capoeira* styles have different techniques and movements speed execution [7]. The aerobic/anaerobic demand imposed to the practitioners [6] seems to be enough to induce chronic cardiovascular adaptations, such as decreased heart rate, and increased markers of autonomic parasympathetic tonus in male beginner practitioners [8].

*Capoeira* training can be organised from strategies with imposed and/or self-selected physical demand (to movements and pace) and rhythmic musical aspects. *Capoeira* progressive training characteristic session can be composed of *capoeira* techniques associated with rhythmic musical elements [4, 8]. The rhythmic musical elements must be produced by the characteristic percussion and development of the typical songs during a *capoeira* training session. The rhythmic element, as well as physical demand on *capoeira* practitioner, still needs to be investigated. However, it is speculated that these elements

can bring an important motivational aspect during the *capoeira* training session because it is self-selected nature.

Little information is available on psychophysiological responses in martial arts and combat sports [9]. To the best of our knowledge, no studies have investigated the physiological and perceptual responses during a single *capoeira* training session. Regarding physiological response, heart rate (HR) is commonly used to assess and monitor physical activity intensity level [10], given its linear relationship with oxygen uptake [11]. As a practical perspective, a rating of perceived exertion (RPE) scale is a simple tool that also can be used to assess and monitor physical activity intensity level [12]. Both HR and RPE have been recommended by the American College of Sports Medicine (ACSM) [13] to prescribe and monitor physical activity intensity, and those moderate-vigorous activities have been recommended to improve fitness and health. Given that the internal load (i.e., stress imposed to the organism) during a *capoeira* training session is unknown it seems important to characterise its intensity using an integrative approach involving an objective (i.e., HR) and a subjective (i.e., RPE) marker of internal load [14].

In addition, to induce a moderate-vigorous internal load to the practitioners, the ACSM [13] has suggested that positive basic affective response (i.e., feeling of pleasure) during exercise/physical activity is a determinant of participation and adherence individual, as previously reported in observational studies [15, 16]. Thus, strategies to enhance the likelihood of gaining pleasurable feelings are likely to contribute to exercise maintenance and subsequent benefits to health and fitness. However, no information is available

about the basic affective response during a single *capoeira* training session. It should be noted that RPE and the basic affective response are not isomorphic constructs. In particular, while the former describes “what” a person feels, the latter emphasises “how” a person feels [17].

In a practical perspective, integrative information considering physiological and perceptual responses during a single *capoeira* training session may contribute to better understand the potential of this modality to improve fitness and health. For example, does *capoeira* meet the physical activity intensity level (i.e., moderate-vigorous) recommended by the ACSM to improve fitness and health in apparently healthy adults [13]? If so, is *capoeira* perceived as a pleasant physical activity by beginners?

The aim was the knowledge about heart rate (HR), rate perceived exertion (RPE) and basic affective responses during different moments of a single *capoeira* progressive training session (CPTS).

We hypothesised that a *capoeira* training session would meet the physical activity intensity level recommended by the ACSM and the participants would report positive basic affective responses during its practice.

## MATERIAL AND METHODS

### Subjects

After signing an informed consent form, a total of 16 physically active adults (10 male and 6 female) were included in this study. The main characteristics of the participants are presented in Table 1. The exclusion criteria were: 1) having any kind of bone, muscle or joint impairment that would preclude participating in the study; 2) having any kind of circulatory or cardiometabolic disease reported in the former health history.

The study was conducted in accordance with the Declaration of Helsinki and was approved by the Research and Ethics Committee of the Federal University of Vale do São Francisco (protocol 0001/200813 CEDEP).

### Experimental design

To test our hypothesis, healthy adults were investigated from fifth visits in the Federal University

of Vale do São Francisco. Aerobic fitness form of anaerobic threshold and maximal power during maximal aerobic incremental exercise test was evaluated. The method used to apply the *capoeira* progressive training session (CPTS) was based on movements of basic programmed lessons. The CPTS was divided into different moments in according to the basic needs of modality. Before and during the CPTS, the participants underwent continuous monitoring of HR and perceptual responses evaluation.

### Aerobic fitness assessment

In a first visit, the participants performed a maximal aerobic incremental exercise test in cycle ergometer (CEFISE®; São Paulo, Brazil). The protocol is starting at 25 watts to female and 50 watts to male and had increments of 25 watts every one min until voluntary exhaustion. The participants maintained 50 RPM's during the entire test. The aerobic fitness indicators analysed were the workload corresponding to the anaerobic threshold (AT) as the aerobic capacity and the workload corresponding to the maximal power ( $P_{MAX}$ ) as the aerobic power [18, 19].

To determination, the AT has adopted the method based on the HR variability threshold in according to Sales et al. [20]. The R-R (see glossary) intervals series was recorded by the HR monitor (*Polar*® model RS800CX, Electro Oy, Kempele, Finland) and filtered in the Polar Precision Performance (v. 4.0) software. This equipment offers high reproducibility and validity [21]. All analyses were run through the HRV Analysis version 2.0 *Kubios* software (Biosignal Laboratory, University of Kuopio, Finland). The HR variability (HRV) was identified through the responses of root mean the successive square difference between adjacent R-R intervals (rMSSD) and standard deviation of instantaneous beat-to-beat R-R interval variability (SD1) by considering the minute of each incremental stage and was named as  $AT_{HRV}$ . During the maximal aerobic incremental exercise test, the  $AT_{HRV}$  was determined when occurred a stabilisation point lower than three milliseconds for the vagal activity indices (SD1 and rMSSD), and the workload of this point was considered. The  $P_{MAX}$  was considered from workload at the moment corresponding to the end stage of the maximal aerobic incremental exercise test that the participant maintained a minimum of the 30s of exercise in according to the established protocol. The  $P_{MAX}$  can be considered as the exercise

**Physical activity** – noun exercise and general movement that a person carries out as part of their day [31].

**Anaerobic threshold** – noun same as the onset of blood lactate accumulation [31].

**Internal** – adjective 1. located within or affecting the inside of something, especially the inside of the body 2. between the members of a team [31].

**Work** verb 1. to exert physical or mental effort in order to do, make or accomplish something, or make someone do this 2. to move or exercise a muscle or part of the body, or be moved or exercised [31].

**Load** – noun 1. a weight or mass which is supported 2. the force that a body part or structure is subjected to when it resists externally applied forces 3. the amount of something, usually weight, that a body part can deal with at one time [31].

**RR** – on the electrocardiography (ECG or EKG), instantaneous heart rate is calculated using the R wave-to-R wave (RR) interval and multiplying/dividing in order to derive heart rate in heartbeats/min. Multiple methods exist:  $HR = 1,500 / (RR \text{ interval in millimetres})$ ;  $HR = 60 / (RR \text{ interval in seconds})$ ;  $HR = 300 / \text{number of "large" squares between successive R waves}$ ;  $HR = 1,500 \text{ number of large blocks} [Wikipedia]$ .

**Cardiorespiratory endurance** – noun the body's ability to carry out prolonged exercise, taking into account both muscle strength and aerobic capacity [31].

intensity corresponding to the maximal oxygen consumption and is the best index to reflect the association between maximum aerobic power and economy of movement [18, 19].

**Capoeira progressive training session (CPTS)**

Previously in distinct days separated by a one week period each, the participants performed three sessions of *progressive capoeira* training to familiarisation with the movements and rhythmic requirement of the modality. The technical improvement naturally occurred every week, resulting in higher speed in movements execution.

In a fifth visit the CPTS experimental protocol, based on the modern styles of *capoeira* [22], as applied by an instructor with 17 years of experience in teaching *capoeira*. The protocol followed the directions of the basic programmed lesson in according to Moreira et al. [8]. This method allows the initiation of novices without any kind of experience in *Capoeira* and can be adapted to individuals in an intermediate and advanced level. The Figure 1 [8] shows the main movements adopted in the CPTS.

The CPTS experimental protocol composition, which lasted for 60 minutes, was from of (a) 2 minutes of playing rhythm instruments – for such an *atabaque* and *pandeiro* were used, which are characteristic rhythmic instruments of *capoeira* with membrafonic design; (b) 2 minutes singing typical songs of *capoeira*; (c) 8 minutes to warm-up with characteristic movements of the modality in a low intensity and static position; (d) 3 sets with 2 minutes of dynamic *ginga* with self-paced nature (main movement of *capoeira* shown in Figure 1 – *ginga<sup>abc</sup>*) by 1 minute of recovery each set; (e) 3 sets with 2 minutes of technical movements interspersed with *ginga* (self-paced nature) following sequence of movements of the Figure 1 (*esquiva lateral* until *esquiva em diagonal*) by 1 minute of recovery each set; (f) 12 minutes with passive participation in *capoeira* circle – at this moment the participants remained in a circle singing characteristics songs of *capoeira* and clapping to the rhythm of the music to watch the *capoeira* game between other two participants (to watch double game); (g) 1 minute of *capoeira* active game in the circle with self-paced and self-selected movements – at this moment the participant



**Figure 1.** The main movements adopted in the basic programmed lesson of *capoeira*. Images from the instructor of *capoeira* progressive training session (adapted from Moreira et al. [8]).

performed freely several movements of *capoeira* (Figure 1) paired with another participant (to perform double game). The remaining time (23 minutes to complete the 60 minutes of the session) was distributed in recovery between the different moments during the CPTS.

### Heart rate response

Ten minutes prior and during the *capoeira* progressive training session the HR was constantly obtained through the Zephyr™ mod. *BioHarness™* 3 monitoring system (Zephyr Technology Corporation, Annapolis, USA). The system enabled the monitoring of all participants during the CPTS, and it consists of a wireless sensor fixed on the chest of each participant by a strap (Zephyr™ mod. ZSFBH). The sensor transmitted the individual information of HR to a signal receiver (Zephyr™ ModFlex GateWay USB) connected to the personal computer and signal repeater (Zephyr™ repeater). Subsequently, the HR responses expressed in “bpm” were analysed moment by moment of CPTS through specific Zephyr™ software. Besides, the HR in the different moments of CPTS was calculated as a percentage of HR during the  $AT_{HRV}$  ( $\%HR-AT_{HRV}$ ) and  $P_{MAX}$  ( $\%HR-P_{MAX}$ ).

### Rating of perceived exertion

RPE was defined as the internal and subjective intensity of effort that was felt during exercise [23]. The whole-body perceived exertion during the CPTS was assessed using the Borg's RPE (6-20) Scale. Before the maximal aerobic incremental exercise test, the meaning of perceived exertion was explained to the subjects. Perceived exertion was defined as the subjective intensity of effort, strain, and/or fatigue that the subjects can feel during exercise [12]. The low and high perceptual anchors for the Borg's RPE scale were established during the maximal aerobic incremental exercise test. A rating of 6 (low anchor, “very, very light”) was assigned to the lowest exercise intensity, while a rating of 20 (high anchor, “very, very hard”) was assigned to the highest exercise intensity.

### Basic affective response

The Feeling Scale (FS) is an 11-point bipolar scale ranging from +5 (plus) to -5 (minus), commonly used to measure the affective response (pleasure/displeasure) during exercise [17]. This scale presents the following verbal anchors: -5 very bad; -3 bad; -1 fairly bad; 0 neutral; +1 fairly

good; +3 good; and +5 very good. The subjects received standard instructions regarding the use of the FS in the initial screening, before the maximal aerobic incremental exercise test, and before the CPTS, according to Hardy and Rejeski [17]. An individual participating in exercise it is quite common to experience changes in mood. Some individuals find exercise pleasurable, whereas others find it to be unpleasurable. Additionally, the feeling may fluctuate across the time. That is, one might feel good and bad a number of times during exercise.

The perceived activation (PA) scale was used to assess the arousal state of the participants [24]. The PA scale is a 6-point single-item measure, ranging from +1 (low arousal) to +6 (high arousal). This scale assesses how the participant felt activated, stimulated and motivated during different times of the study (before and during the CPTS). The affective valence measured by the FS is a fundamental component of the circumplex model of Russell [25], which incorporates affective valence and PA as an orthogonal and bipolar dimension of the affective space.

Finally, standard definitions of perceptual responses and separate instructional sets for scales were read to the participants immediately before the maximal aerobic incremental exercise test. The low and high perceptual anchors for the RPE, FS and PA scale were established during the maximal aerobic incremental exercise test. RPE, FS and PA values were evaluated randomly during the last 10 seconds of each stage of maximal aerobic incremental exercise test and immediately after the different moments of CPTS (after 10 minutes of resting and session moments).

### Statistical analysis

Data are expressed as the mean and standard deviation (SD or  $\pm$ ). Data were tested for normality using the Shapiro-Wilk test. One-way repeated measures ANOVA reporting F-ratio, degrees of freedom and P-value were used to verify the main effects of time in the different moments of CPTS. Mauchly's test was analysed by assessing the data sphericity. In the case of sphericity assumption violation, the degrees of freedom were adjusted and reported using the Greenhouse-Geisser adjustment. Partial eta squared ( $\eta_p^2$ ) was used to determine the effect size. When the interaction of time was found,



Bonferroni-corrected multiple pairwise comparisons and adjusted P-values were reported. Student's t-test for paired samples was performed to verify the effects of time from resting to *capoeira* total session results. Student's t-test for independent samples was performed to compare the general characteristics between male and female groups. As a result of the t-tests were reported t-value, 95% confidence interval and P-value. The level of significance adopted was set at  $p < 0.05$  and the software used for analysis was the SPSS 22.0 for Windows (SPSS, Inc., Chicago, USA).

## RESULTS

Males presented higher values of  $AT_{HRV}$  ( $t = 4.838$  [CI (confidence interval):  $-94.54, -36.46$ ];  $p < 0.001$ ) and  $P_{MAX}$  ( $t = 5.544$  [CI:  $-130.60, -57.73$ ];  $p < 0.001$ ) than females. However, males and females had similar age ( $t = 1.132$  [CI:  $-3.51, 11.38$ ];  $p = 0.276$ ), body mass ( $t = 1.510$  [CI:  $-26.85, 4.66$ ];  $p = 0.153$ ), body mass index ( $t = 0.208$  [CI:  $-4.43, 5.39$ ];  $p = 0.838$ ), and physical activity level ( $t = 0.556$  [CI:  $-204.75, 119.75$ ];  $p = 0.586$ ). The results of internal loads corresponding to aerobic fitness did not differ between the sexes, as noted in  $AT_{HRV}$

**Table 1.** Mean and standard deviation ( $\pm$ ) of descriptive characteristics of the sample and aerobic fitness responses during the maximal aerobic incremental test.

Variable (indicator)	Total (n = 16)	Male (n = 10)	Female (n = 6)
<b>General characteristics</b>			
Age (years)	27.9 $\pm$ 6.8	26.4 $\pm$ 5.2	30.3 $\pm$ 8.8
Body mass (kg)	75.3 $\pm$ 14.8	79.5 $\pm$ 12.8	68.4 $\pm$ 16.5
Height (cm)	170.0 $\pm$ 0.9	175.0 $\pm$ 0.4	<b>160.0 <math>\pm</math> 0.5*</b>
Body mass index ( $\text{kg}\cdot\text{m}^{-2(-1)}$ )	26.1 $\pm$ 4.3	25.9 $\pm$ 4.0	26.4 $\pm$ 5.1
Physical activity ( $\text{min}\cdot\text{week}^{-1}$ )	306.3 $\pm$ 142.6	322.0 $\pm$ 119.3	280.0 $\pm$ 184.6
<b><math>AT_{HRV}</math></b>			
Power (W)	141 $\pm$ 41	166 $\pm$ 31	<b>100 <math>\pm</math> 14***</b>
Heart rate (bpm)	146 $\pm$ 10	144 $\pm$ 12	149 $\pm$ 8
RPE (rate perceived exertion)	12 $\pm$ 3	12 $\pm$ 3	12 $\pm$ 2
FS (Feeling Scale)	1 $\pm$ 3	1 $\pm$ 3	1 $\pm$ 3
PA (perceived activation)	4 $\pm$ 1	4 $\pm$ 2	4 $\pm$ 1
<b><math>P_{MAX}</math></b>			
Power (W)	248 $\pm$ 60	278 $\pm$ 36	<b>183 <math>\pm</math> 26***</b>
Heart rate (bpm)	181 $\pm$ 12	179 $\pm$ 14	183 $\pm$ 6
RPE (rate perceived exertion)	19 $\pm$ 2	19 $\pm$ 2	19 $\pm$ 2
FS (feeling scale)	-2 $\pm$ 4	-1 $\pm$ 4	-3 $\pm$ 4
PA (perceived activation)	6 $\pm$ 1	6 $\pm$ 1	6 $\pm$ 1

$AT_{HRV}$  anaerobic threshold measured by heart rate variability during the maximal aerobic incremental test;  $P_{MAX}$  maximal power obtained during the maximal aerobic incremental test; \* $p < 0.05$  and \*\*\* $p < 0.001$  to male.

to HR (t = 0.834 [CI: -7.11, 16.18]; p=0.418), RPE (t = 0.620 [CI: -3.57, 2.20]; p=0.620), FS (t = 0.256 [CI: -3.59, 2.82]; p=0.801) and PA (t = 0.720 [CI: -1.92, 0.95]; p=0.482) and in P<sub>MAX</sub> to HR (t = 0.654 [CI: -9.03, 16.96]; p=0.523), RPE (t = 0.720 [CI: -2.38, 1.18]; p=0.483), FS (t = 0.510 [CI: -5.72, 3.52]; p=0.617) and PA (t = 0.366 [CI: -1.36, 0.96]; p=0.719) (Table 1).

The similarities between the sexes concern of the internal loads responses corresponding to the aerobic fitness, the absolute HR and relative HR to AT<sub>HRV</sub> and P<sub>MAX</sub> in relation to the different moments of *capoeira* progressive training session (Table 2). When compared to resting was found a main effect of time to absolute HR (F<sub>[3,262,48,926]</sub> = 53.107; p<0.0001; η<sub>p</sub><sup>2</sup> = 0.780) and relative HR to AT<sub>HRV</sub> (F<sub>[3,407,51,099]</sub> = 50.430; p<0.0001; η<sub>p</sub><sup>2</sup> = 0.771) and P<sub>MAX</sub> (F<sub>[3,393,50,893]</sub> = 51.011, p<0.0001; η<sub>p</sub><sup>2</sup> = 0.773).


There was a main effect of time for RPE (F<sub>[2,499,37,486]</sub> = 45.774; p<0.001; η<sub>p</sub><sup>2</sup> = 0.753), whereas the FS did not change between the different moments of CPTS (F<sub>[2,101,31,509]</sub> = 1.703; p=0.197; η<sub>p</sub><sup>2</sup> = 0.102). Also, a main effect of time was observed for PA (F<sub>[2,105,31,570]</sub> = 15.857; P=0.001; η<sub>p</sub><sup>2</sup> = 0.514) with results in the resting (3 ± 1 points), statistically different (p<0.01) to moments singing songs (4 ± 1 points), *ginga* (4 ± 1 points), technical movements (5 ± 1 points) and *capoeira* active game (5 ± 1 points) (Figure 2).

When compared to resting was found statistically differences to absolute HR (t = -21.447 [CI: -43.21, -35.40]; p<0.0001) and relative HR to AT<sub>HRV</sub> (t = -21.830 [CI: -29.49, -24.25]; p<0.0001) and P<sub>MAX</sub> (t = -19.509 [CI: -24.26, -19.48]; p<0.0001). Besides, the Figure 3D and 3F show that the perceptual variables also increased in CPTS total when analyzed the

**Table 2.** Mean and standard deviation (±) of absolute heart rate (HR) in different moments of *capoeira* progressive training session and corresponding HR to anaerobic threshold (%HR-AT<sub>HRV</sub>) and maximal power (%HR-P<sub>MAX</sub>) during the maximal aerobic incremental test (n =16).

Variable (indicator)	HR (bpm)	HR-AT <sub>HRV</sub> (%)	HR-P <sub>MAX</sub> (%)
Resting (10min)	77 ±13	53 ±10	43 ±7
<b>Capoeira progressive training session</b>			
Playing instruments (2min)	87 ±13	60 ±9	48 ±7
Recovery (3min)	<b>88 ±10***</b>	<b>60 ±8***</b>	<b>49 ±5***</b>
Singing songs (2min)	84 ±14	58 ±10	47 ±7
Recovery (1min)	<b>88 ±11***</b>	<b>61 ±9***</b>	<b>49 ±6***</b>
Warm-up (8min)	<b>102 ±12***</b>	<b>70 ±1***</b>	<b>57 ±8***</b>
Recovery (2min)	<b>104 ±15***</b>	<b>72 ±1***</b>	<b>58 ±9***</b>
<i>Ginga</i> (8min)	<b>124 ±17***</b>	<b>85 ±13***</b>	<b>69 ±10***</b>
Recovery (6min)	<b>102 ±14***</b>	<b>71 ±12***</b>	<b>57 ±8***</b>
Technical movements (8min)	<b>134 ± 5***</b>	<b>92 ±12***</b>	<b>74 ±8***</b>
Recovery (7min)	<b>122 ±16***</b>	<b>84 ±13***</b>	<b>67 ±8***</b>
<i>Capoeira</i> passive game (6min)	<b>118 ±25***</b>	<b>80 ±15***</b>	<b>65 ±11***</b>
<i>Capoeira</i> active game (1min)	<b>135 ±25***</b>	<b>93 ±18***</b>	<b>75 ±13***</b>
<i>Capoeira</i> passive game (6min)	<b>129 ±17***</b>	<b>89 ±16***</b>	<b>72 ±10***</b>

\*\*\*p<0.0001 to resting (adjusted p values by Bonferroni-corrected multiple pairwise comparisons).



**Figure 2.** Mean and standard error (SE) of rate perceived exertion (RPE) and feeling scale (FS) responses during the *capoeira* progressive training session and corresponding value to anaerobic threshold ( $AT_{HRV}$ ) during the maximal aerobic incremental test ( $n = 16$ ). \* $p < 0.01$  to resting (adjusted P-values by Bonferroni-corrected multiple pairwise comparisons).

RPE ( $t = -5.984$  [CI:  $-4.84, -2.30$ ];  $p < 0.0001$ ) and PA ( $t = -5.238$  [CI:  $-2.11, -0.88$ ];  $p < 0.0001$ ), whereas the FS (Figure 3E) remained unchanged from resting to CPTS total ( $t = 0.211$  [CI:  $-1.02, 1.24$ ];  $p = 0.836$ ) (Figure 3).

## DISCUSSION

Our initial hypothesis was confirmed. We highlight that this is the first study that used an integrative psychophysiological approach to characterise a *capoeira* progressive training session (CPTS).


Our results suggest that a regular *capoeira* training program may be an alternative form of physical activity to improve fitness and health. Three studies [6-8] involving *capoeira* exercise and HR responses were found in the literature. However, none of them presented acute responses during the CPTS in male practitioners. The CPTS elicited a mean of  $65 \pm 7\%$  of maximum HR ( $\%HR-P_{MAX}$ ) obtained in the maximal aerobic incremental exercise test (Figure 3-C) and  $11 \pm 2$  in the 15-point Borg scale (Figure 3-D). Also, during the session occurred a change between  $47 \pm 7\%$  and  $75 \pm 13\%$  of maximum HR of the sample (Table 2) and RPE between  $8 \pm 2$  and  $14 \pm 2$  in the 15-point Borg scale

(Figure 2). The ACSM from its position stand [13] suggests a classification of exercise intensity for cardiorespiratory endurance, where moderate exercise intensity is represented by 64-76% of maximum HR or RPE between 12 and 13 in the 15-point Borg scale. In this way, it is possible that when frequently performed, a CPTS meets the assumptions to moderate exercise intensity and subsequently increasing physical fitness in apparently healthy adults [13]. However, a longitudinal study for this purpose needs to be developed.

Specifically, when analysed the RPE as a method to mark the exercise effort [12] in the different moments of CPTS, was possible to note a mean value to *ginga* at  $12 \pm 2$  and, technical movements and *active capoeira* game at  $14 \pm 2$  in the 15-point Borg scale (Figure 2). According to Garber et al. [13], the high-intensity effort (somewhat hard to very hard) is scheduled from 14 to 17 on the 15-point Borg scale. These findings, in addition to demonstrating the intermittent nature of CPTS, also suggests that from investigated moments the application of interval training models in the *capoeira* is possible (high intensity vs moderate intensity), and the use of the Borg scale can be a useful tool in order to control the training loads.

Regarding the basic affective responses, the mean





**Figure 3.** Mean and standard deviation ( $\pm$ ) of absolute heart rate (HR), relative HR to anaerobic threshold ( $\%HR-AT_{HRV}$ ) and maximal power ( $\%HR-P_{MAX}$ ) during maximal aerobic incremental test, rate perceived exertion (RPE), feeling scale (FS) and perceived activation (PA) responses of *capoeira* progressive training total session (n =16): \* $p < 0.0001$  to resting.

value of the session was  $3 \pm 2$  (Figure 3), and values during the technical movements and *active capoeira* game were  $2 \pm 3$  (Figure 2). These values characterise a positive affective response during *capoeira* exercise (i.e., feeling of pleasure) without statistical difference compared to resting ( $p > 0.05$ ; Figures 2 and 3). These findings challenge the traditional model highlighted in the literature that shows that when exercise intensity increases, especially above to the ventilatory threshold, the basic affective response decreases gradually to negative values (i.e., feeling of displeasure). Ekkekakis [27] also suggested that the basic affect may decline due to both the high intensity perceived effort and the loss of autonomy inherent in an imposed exercise prescription. The CPTS was composed by rhythmic musical elements associated with *capoeira* techniques from self-paced nature movements (i.e., *ginga* and technical

movements) and a moment with self-selected and self-paced movements (*active capoeira* game) as demonstrated by Moreira et al. [8]. In this last moment, the practitioners performed freely the *capoeira* movements (Figure 1) in a double game, whereas in the *ginga* and technical movements the practitioners performed such movements within a possible physical condition to your level (self-paced nature). Moreover, the game character/body challenge involved in the CPTS may be contributing to positive responses from basic affect [28], which might reduce the discomfort caused by the intensity of effort [29]. It is speculated that the highlighted characteristics of CPTS maintained positive affective response despite increased effort imposed to the practitioners in different moments of the session (Figure 2). From this perspective, it highlights the results obtained with the PA during the CPTS, where it was found

in the total session ( $4 \pm 1$ ; Figure 3), as well as *ginga* ( $4 \pm 1$ ), technical movements ( $5 \pm 1$ ) and *capoeira* active game ( $5 \pm 1$ ) higher values when compared ( $p < 0.01$ ) to the resting ( $3 \pm 1$ ). These findings shows a higher arousal and/or motivation state of practitioners [23] during these moments of the CPTS, which together with affective valence (Figures 2 and 3) integrating the dimension of the affective space [25] and suggest the *capoeira* as a possible hedonic model of physical activity prescription, which aims to adherence of an individual to the regular physical exercise program [30]. However, a novel longitudinal study for this purpose needs to be developed.

From a practical perspective, the results support the fitness and conditioning professionals with the possibility of using CPTS to improve individual's fitness and health and maintain a long-term adherence. The CPTS prescription would be done according to the basic programmed lesson method (Figure 1) [8] and composed by 60 minutes with musical rhythmic elements (playing rhythm instruments as *pandeiro* and *atabaque* and singing typical songs of *capoeira*) associated with *capoeira* techniques from of movements within a physical condition level of practitioners (self-paced movements and moments with self-selected movements). However, throughout the possible training program, the CPTS may be gradually intensified through the natural gains in the velocity of movements that occurs in parallel to the improvement of technique.

Importantly, the present study presented a limitation related the lack of a measure of maximum

HR during the *capoeira* exercise, which could contribute to greater precision in relation to percentage demand of CPTS. Accordingly, the lack of protocols in the literature that enables the assessment of maximum stress in *capoeira*, the maximum HR was obtained through a maximal aerobic incremental exercise test in cycle ergometer. Although not a specific test, enabled real and not estimates results of maximum HR, which make us believe on the basis of an estimate by the age of the sample that is very close to the possible values in *capoeira* (Table 1). However, a study suggesting a specific protocol within *capoeira* needs to be performed.

## CONCLUSIONS

We concluded that a single CPTS increased the HR, RPE and PA responses significantly, maintaining affective valence (FS) even with increased intensity in the practitioners during the session. Besides, a CPTS may be an alternative physical activity modality to improve fitness and health of adults given that it meets the ACSM [10] recommendation regarding physical activity intensity level (i.e., moderate-vigorous); moreover, participants perceived it as a pleasant activity, which may enhance long-term adherence.

## ACKNOWLEDGEMENTS

The authors thank Iraíma Lago and Karoline Andrade for photo design.

## REFERENCES

1. Maia RB, Martins MCC, Rocha CHL et al. Efeito da Prática de Capoeira sobre os Parâmetros Cardiovasculares. *Rev Bras Cardiol* 2010; 23(1): 68-73 [in Portuguese]
2. Cunha IMCF, Vieira LR, Tavares LCV et al. Capoeira: a memória social construída por meio do corpo. *Rev Bras Cienc Mov* 2014; 22: 735-55 [in Portuguese]
3. Vieira LP, Assunção MR. Os desafios contemporâneos da Capoeira. Brasília: Ministério das Relações Exteriores. *Revista Textos do Brasil* 2009; 9-19 [in Portuguese]
4. Santos AO. "Mestre Mestrinho" Capoeira Arte-Luta Brasileira. Cascavel: Editora Assoeste; 2001 [in Portuguese]
5. Assunção MR. Capoeira: The History of an Afro-Brazilian Martial Art. London, New York: Taylor & Francis; 2005
6. Silva FF, Souza RA, Costa RP et al. Heart rate responses during and after the practice of Capoeira: A Brazilian art form - Part II. *J Exerc Physiol Online* 2013; 16(2): 112-116
7. Silva FF, Souza RA, Carvalho WRG et al. Analysis of acute cardiovascular responses in experienced practitioners of Capoeira: A Brazilian art form. *J Exerc Physiol Online* 2012; 15: 112-119
8. Moreira SR, Teixeira-Araujo AA, Oliveira dos Santos A et al. Ten weeks of Capoeira progressive training improved cardiovascular parameters in male practitioners. *J Sports Med Phys Fitness* 2017; 57(3): 289-98
9. Branco BHM, Andreato LV, Mendes AA et al. Effects of a Brazilian jiu-jitsu training session on physiological, biochemical, hormonal and perceptible responses. *Arch Budo Sci Martial Art Extreme Sport* 2016; 12: 145-54
10. American College of Sports Medicine. ACSM's Guidelines for Exercise Testing and Prescription. 7th ed. Rio de Janeiro: Guanabara Koogan; 2007
11. Moon JK, Butte NF. Combined heart rate and activity improve estimates of oxygen consumption and carbon dioxide production rates. *J Appl Physiol* 1996; 81(4): 1754-1761
12. Borg G. Ratings of perceived exertion and heart rates during short-term cycle exercise and their use in a new cycling strength test. *Inter J Sports Med* 1982; 3(3): 153-158

13. Garber CE, Blissmer B, Deschenes MR et al. American College of Sports Medicine. American College of Sports Medicine Position Stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011; 43(7): 1334-1359
14. Costa EC, Vieira CM, Moreira A et al. Monitoring external and internal loads of Brazilian soccer referees during official matches. *J Sports Sci Med* 2013; 12(3): 559-564
15. Williams DM, Dunsinger S, Ciccolo JT et al. Acute affective response to a moderate-intensity exercise stimulus predicts physical activity participation and 6 and 12 months later. *Psych Sport Exerc* 2008; 9(3): 231-245
16. Williams DM, Dunsinger S, Jennings EG et al. Does affective valence during and immediately following a 10-min walk predict concurrent and future physical activity? *Annals Behav Med* 2012; 44(1): 43-51
17. Hardy CJ, Rejeski WJ. Not what, but how one feels: the measurement of affect during exercise. *J Sports Exerc Psychol* 1989; 11: 304-317
18. Caputo F, Stella SG, Mello MT et al. Indexes of power and aerobic capacity obtained in cycle ergometry and treadmill running: comparisons between sedentary, runners, cyclist and triathletes. *Rev Bras Med Esp* 2003; 9: 231-237
19. Caputo F, Oliveira MFM, Greco CC et al. Aerobic exercise: bioenergetics, physiological adjustments, fatigue and performance indices. *Rev Bras Cineantropom Desempenho Hum* 2009; 11(1): 94-102
20. Sales MM, Campbell CSG, Morais PK et al. Noninvasive method to estimate anaerobic threshold in individuals with type 2 diabetes. *Diabetol Metab Syndr* 2011; 3: 1-8
21. Williams DP, Jarczok MN, Ellis RJ et al. Two-week test-retest reliability of the Polar® RS800CX™ to record heart rate variability. *Clin Physiol Funct Imaging* 2017; 37(6): 776-781
22. Assunção MR. Capoeira circle or sports academy? The emergence of modern styles of capoeira and their global context. *Hist Cienc Saude Manguinhos* 2014; 21(1): 135-149
23. Noble BJ, Robertson RJ. *Perceived Exertion*. Champaign, IL: Human Kinetics; 1996
24. Sveback S, Murgatroyd S. Metamotivational dominance: a multimethod validation of reversal theory constructs. *J Percep Soc Psych* 1985; 48(1): 107-116
25. Russell JA. A circumplex model of affect. *J Pers Soc Psychol* 1980; 39(6): 1161-1178
26. Backhouse SH, Ekkekakis P, Bidle SJ et al. Exercise makes people feel better but people are inactive: paradox or artefact? *J Sport Exerc Psychol* 2007; 29(4): 498-517
27. Ekkekakis P. Let them roam free? Physiological and psychological evidence for the potential of self-selected exercise intensity in public health. *Sports Med* 2009; 39(10): 857-888
28. Lieberman DE. Is Exercise Really Medicine? An Evolutionary Perspective. *Curr Sports Med Rep* 2015; 14(4): 313-319
29. Lee HH, Emerson JA, Williams DM. The Exercise-Affect-Adherence Pathway: An Evolutionary Perspective. *Front Psychol* 2016; 7: 1285
30. Jung ME, Bourne JE, Little JP. Where does HIT fit? An examination of the affective response to high-intensity intervals in comparison to continuous moderate- and continuous vigorous-intensity exercise in the exercise intensity-affect continuum. *PLoS One* 2014; 9: e114541
31. *Dictionary of Sport and Exercise Science. Over 5,000 Terms Clearly Defined*. London: A & B Black; 2006

**Cite this article as:** Moreira SR, Teixeira-Araujo AA, Camara MVF et al. Heart rate, a rating of perceived exertion and basic affective responses during different moments of a single capoeira progressive training session (CPTS). *Arch Budo Sci Martial Art Extreme Sport* 2017; 13: 153-163